

My NCBI
[Sign In] [Re

[All Databases](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[OMIM](#)[PMC](#)[Journals](#)

Search **PubMed** for **weinstein telomere** [Go](#) [Clear](#) [Save S](#)

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)

Display **Summary** Show **20** Sort by Send to

[About Entrez](#)[Text Version](#)

Entrez PubMed

[Overview](#)[Help](#) [FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)

PubMed Services

[Journals Database](#)[MeSH Database](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[Special Queries](#)[LinkOut](#)[My NCBI](#)

Related Resources

[Order Documents](#)[NLM Mobile](#)[NLM Catalog](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)

☐ **1:** [Weinstein BS, Ciszek D.](#)

[Related Articles, Links](#)



The reserve-capacity hypothesis: evolutionary origins and modern implications of the trade-off between tumor-suppression and tissue-repair.

Exp Gerontol. 2002 May;37(5):615-27. Review.

PMID: 11909679 [PubMed - indexed for MEDLINE]

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Nov 3 2005 04:38:43



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show 20 Sort by Send to

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Curr Mol Med. 2005 Mar;5(2):197-203.

Related Articles, Links

Telomeres as biomarkers for ageing and age-related diseases.

von Zglinicki T, Martin-Ruiz CM.

Henry Wellcome Laboratory for Biogerontology Research, Newcastle General Hospital, School of Clinical Medical Sciences-Gerontology, University of Newcastle, Newcastle upon Tyne, NE4 6BE, UK.

Telomeres in telomerase-negative cells shorten during DNA replication in vitro due to numerous causes including the inability of DNA polymerases to fully copy the lagging strand, DNA end processing and random damage, often caused by oxidative stress. Short telomeres activate replicative senescence, an irreversible cell cycle arrest. Thus, telomere length is an indicator of replicative history, of the probability of cell senescence, and of the cumulative history of oxidative stress. Telomeres in most human cells shorten during ageing in vivo as well, suggesting that telomere length could be a biomarker of ageing and age-related morbidity. There are two distinct possibilities: First, in a tissue-specific fashion, short telomeres might indicate senescence of (stem) cells, and this might contribute to age-related functional attenuation in this tissue. Second, short telomeres in one tissue might cause systemic effects or might simply indicate a history of high stress and damage in the individual and could thus act as risk markers for age-related disease residing in a completely different tissue. In recent years, data have been published to support both approaches, and we will review these. While they together paint a fairly promising picture, it needs to be pointed out that until now most of the evidence is correlative, that much of it comes from underpowered studies, and that causal evidence for essential pathways, for instance for the impact of cell senescence on tissue ageing in vivo, is still very weak.

Publication Types:

- [Review](#)
- [Review, Tutorial](#)

PMID: 15974873 [PubMed - indexed for MEDLINE]